

The seasonal investigation of Symbiodiniaceae in broadcast spawning, *Acropora humilis* and brooding, *Pocillopora* cf. *damicornis* corals

Suppakarn Jandang¹, Voranop Viyakarn^{1,2}, Yuki Yoshioka³, Chuya Shinzato³ and Suchana Chavanich^{1,2,4}

¹ Reef Biology Research Group, Department of Marine Science, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

² Aquatic Resources Research Institute, Chulalongkorn University, Bangkok, Thailand

³ Atmosphere and Ocean Research Institute, The University of Tokyo, Chiba, Japan

⁴ Center of Excellence for Marine Biotechnology, Department of Marine Science, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

ABSTRACT

The density and diversity of Symbiodiniaceae associated with corals can be influenced by seasonal changes. This study provided the first annual investigation of Symbiodiniaceae density and diversity associated with *Acropora humilis* and *Pocillopora* cf. *damicornis* corals in the Gulf of Thailand using both zooxanthellae cell count and next-generation sequencing (ITS-1, ITS-2 regions) techniques, respectively. The results from this study indicated that zooxanthellae cell densities in both coral species differ significantly. The number of zooxanthellae was negatively correlated with the physical environment variable (light intensity). The diversity within *A. humilis* consisted of two genera, *Cladocopium* (Cspc_C3: 56.39%, C3w: 33.62%, C93type1: 4.42% and Cspf: 3.59%) and a small amount of *Durusdinium* (D1: 1.03%) whereas *P. cf. damicornis* was found to be 100% associated with *Durusdinium* (D1: 95.58%, D6: 1.01% and D10: 2.7%) suggesting that each coral species may select their appropriate genus/species of Symbiodiniaceae in response to local environmental stressors. The results of this study provided some information on the coral-Symbiodiniaceae relationship between seasons, which may be applied to predict the potential adaptation of corals in localized reef environments.

Subjects Biodiversity, Ecology, Genetics, Genomics, Marine Biology

Keywords Zooxanthellae, *Cladocopium*, *Durusdinium*, Next Generation Sequencing, Corals, Seasonal, Thailand, Symbiodiniaceae, *Acropora*, *Pocillopora*

INTRODUCTION

Coral reefs act as habitats of high ecological complexity and biodiversity in the marine environment (*Blackall, Wilson & van Oppen, 2015; Brown et al., 1999a*). Most reef-building corals are known to sustain a symbiotic relationship with dinoflagellate algae belonging to the family Symbiodiniaceae (also known colloquially as zooxanthallae) (*Eckert et al., 2020; LaJeunesse, 2002*). Nutrient exchange between coral and Symbiodiniaceae underpin the success of reef ecosystems in oligotrophic waters (*Ferrier-Pagès et al., 1998; Pierangelini, Thiry & Cardol, 2020*). The photosynthetic products (carbon (C), nitrogen (N), and

Submitted 15 September 2021

Accepted 23 February 2022

Published 14 June 2022

Corresponding author

Suchana Chavanich,
Suchana.c@chula.ac.th

Academic editor
Anastazia Banaszak

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.13114

© Copyright
2022 Jandang et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

phosphorus (P)) supplied by Symbiodiniaceae are the primary means of support to host productivity, metabolism, growth and reproduction while the host coral provides the needed compounds to their symbiont for photosynthesis (Peng et al., 2012). Therefore, the symbiosis between corals and Symbiodiniaceae is essential for the development and survival of coral reefs ecosystems (Baker et al., 2013; Wilkinson et al., 2015). The density and diversity of Symbiodiniaceae associated with corals is influenced by different factors including ecological and physical conditions (Terraneo et al., 2019; Thornhill et al., 2009). Seasonal variation of the coral-Symbiodiniaceae relationship is known to effect coral health indices such as Chlorophyll *a* and zooxanthellae density, due to both physical and chemical factors of the surrounding environment (Hinrichs et al., 2013; Scheufen, Iglesias-Prieto & Enríquez, 2017). The density of zooxanthellae in several scleractinian corals around the world has been shown to be variable throughout the year, in response to seasonality (Li et al., 2008; Stimson, 1997; Zhou et al., 2017). High irradiance levels and temperatures may have severe impacts on zooxanthellae density (Fisher, Malme & Dove, 2012; Glynn, 1993; Pillay, Willis & Terashima, 2005). Most tropical reef-building corals possess their highest zooxanthellae densities during winter months, declining in the summer (Brown et al., 1999a; Fitt et al., 2001; Stimson, 1997). This has been shown to correlate with minimal coral growth rates, which occur simultaneously with low zooxanthellae density during summer season (Al-Hammady, 2013; Pillay, Willis & Terashima, 2005).

Reef-building corals possess symbiotic relationships with a genetically diverse (genera/species) range of Symbiodiniaceae, which have been attributed to coping with different environmental stressors (Rouze et al., 2019; Silverstein et al., 2011; Silverstein, Correa & Baker, 2012). However, the revised systematics of zooxanthellae were recently defined based on genetic variation (LaJeunesse et al., 2018). Seven genera of Symbiodiniaceae were described: *Symbiodinium*, *Breviolum*, *Cladocopium*, *Durusdinium*, *Effrenium*, *Fugacium* and *Greakladium* (formerly clades A, B, C, D, E, F and G, respectively). The physiological responses of corals to environmental stressors may differ among different genera/species of Symbiodiniaceae (DeSalvo et al., 2010; Rowan, 2004; Sampayo et al., 2008). The selective specific uptake of symbiont species could lead to differentiation of coral fitness and survival rates (Quigley, Willis & Bay, 2016; Tan et al., 2020). Corals associated with *Durusdinium* (formerly clade D) exhibit less damage to photosystem II (PSII) compared to others mutualistic Symbiodiniaceae genera (Jones & Berkelmans, 2011; Little, Van Oppen & Willis, 2004; Suwa, Hirose & Hidaka, 2008). Conversely, corals predominantly containing the genus *Cladocopium* (formerly clade C) had translocated large amounts of fixed carbon to their host, which could affect its capacity to remain metabolically active under high temperature conditions (Cantin et al., 2009).

To minimize the impact on coral performance, most corals do not change their Symbiodiniaceae community structure (Cunning, Silverstein & Baker, 2018; Goulet, 2006). However, the ability of corals to change symbiont proportion and/or species has consistently correlated with disturbance severity and the diversity of Symbiodiniaceae species within the host (Cunning, Silverstein & Baker, 2015; Rouze et al., 2019). Some corals adapt or acclimatize when living in more hostile environments *via* two potential adaptative mechanisms, namely, “shuffling” (change in relative abundance of Symbiodiniaceae species

within a colony) and “switching” (uptake of new exogenous Symbiodiniaceae species from the environment) ([Berkelmans & Van Oppen, 2006](#); [Rouze et al., 2016](#); [Sampayo et al., 2008](#)). Several studies have corroborated that the genus *Durusdinium* may be found at significantly higher proportions during thermal stress than normal environmental conditions ([Chen et al., 2020](#); [Cunning, Silverstein & Baker, 2015](#); [Dilworth et al., 2021](#); [Poquita-Du et al., 2020](#); [Stat et al., 2013](#)). Moreover, the increase of opportunistic *Durusdinium* cells may help corals survive under thermal stress by assimilating and transferring the reduced amounts of organic compounds to the coral host ([Fabricius et al., 2004](#); [Ghavam Mostafavi et al., 2007](#); [Wall et al., 2020](#)). Consequently, bleached corals may become more resistant to higher temperatures by symbiont shuffling and/or switching ([Császár et al., 2010](#); [Davies et al., 2018](#)). As such, the flexibility of the coral-Symbiodiniaceae relationship is important in understanding their future survival in response to environmental stressors in particular areas and coral species ([Baskett, Gaines & Nisbet, 2009](#); [Sampayo et al., 2016](#)).

The different reproductive modes of corals can affect the establishment of the coral endosymbiosis. Generally, broadcast spawning corals acquire their Symbiodiniaceae from the surrounding environment (horizontal symbiont transmission: HT), whereas in contrast, brooding corals transfer Symbiodiniaceae cells from parents to offspring (vertical symbiont transmission: VT) ([Baird et al., 2009](#); [Sweet, 2014](#)). In general, most corals that use horizontal symbiont acquisition are able to select a genetically diverse assemblage of symbiont partners, conferring a greater opportunity to shift their dominant Symbiodiniaceae composition, especially when corals face unstable environmental conditions ([Boilard et al., 2020](#); [Thornhill et al., 2017](#)). Conversely, corals that use vertical symbiont transmission are theoretically able to harbor stable communities of Symbiodiniaceae due to genetic regulation of brooding corals ([Quigley et al., 2018](#); [Thornhill, Fitt & Schmidt, 2006](#)). Therefore, Symbiodiniaceae specificity, and the stability of communities, are predicted to be improved when symbionts are transmitted vertically compared to horizontally ([Baker, 2003](#); [Thornhill, Fitt & Schmidt, 2006](#)). However, recent studies revealed that brooding corals (VT) become more flexible by acquiring resilient species of Symbiodiniaceae from the environment (HT) and are able to transfer the new Symbiodiniaceae communities to their offspring ([Quigley et al., 2018](#); [Quigley, Willis & Kenkel, 2019](#)).

In Thailand, almost all Acroporidae are believed to be HT species while some species of Pocilloporidae including *P. cf. damicornis* are VT species ([Kuanui et al., 2008](#)). Little is known about the variability and dynamics between Symbiodiniaceae and their coral hosts particularly in Thai waters. Thus, this study provides the first investigation of Symbiodiniaceae communities and diversity associated with different reproductive modes corals: *Acropora humilis* (a gonochoric broadcast spawner: HT) and *Pocillopora cf. damicornis* (a hermaphroditic brooder: VT) from the upper Gulf of Thailand over a one year period.

MATERIALS & METHODS

Sampling site and sample collection

Ko Tao Mo (KTM) is located at the Royal Thai Navy Base, Chon Buri province, upper Gulf of Thailand (GoT) ($12^{\circ}38'33.78''\text{N}$ $100^{\circ}51'40.13''\text{E}$) (Fig. 1). The average depth of the sea floor is approximately 3–15 m. Most coral communities are mainly found around the inner area of the breakwater next to KTM, after it was created in 1995. Therefore, this new coral community was estimated to be 23 years old at the time of sampling. A total of 67 species of reef-building corals were recorded. *Acropora* was found as a dominant genus in this area (Viyakarn *et al.*, 2009). In this study, *Acropora humilis* and *Pocillopora cf. damicornis* corals were collected around KTM. These corals were distributed in shallow water at 5–7 m depth. Coral colonies ($n = 15$) were tagged for each species. The minimum distance between each selected colony was 3 m apart to maximize the chance of sampling distinct genetically diverse of corals. The size of tagged colonies ranged from 30–100 cm in diameter. Coral samples were collected following different seasons, referring to the local information of the Thailand Meteorological Department. These were divided into three seasons including winter (November–February), summer (March–June) and rainy (July–October). Samples were collected in January (winter), May (summer) and October (rainy) 2018, respectively. Fragments of approximately 2–3 cm from tagged colonies of each coral species and season within a year were collected by SCUBA diving. In addition, ambient sea water temperature and light intensity data were recorded using underwater temperature data loggers (Hobo, Onset Corporation Ltd, Bourne, MA, USA). The average light intensity and temperature of each season were calculated based on logged data from the winter season of 2017 until the rainy season in 2018. The average temperature and light intensity (lux) was converted to PAR, $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ (approximately $1 \mu\text{mol m}^{-2} \text{s}^{-1} = 52 \text{ lux}$) following the calculation method according to Bednarz *et al.* (2015).

Determination of zooxanthellae density

Fresh coral fragments were collected seasonally from each of the 15 tagged colonies of *A. humilis* and *P. cf. damicornis* corals using hammer and chisel. Coral fragments were then transferred to laboratory and preserved in 10% buffered formalin. For zooxanthellae density determination, we adapted the decalcification method from McCowan *et al.* (2011). Briefly, preserved coral branches were placed in a glass container with 5% HCl solution until the coral skeleton was completely dissolved (~ 4 –7 days depending on size and species). The remaining tissues were rinsed and preserved in 70% ethanol. Three pieces of fixed tissue were cut into 1×1 cm sections and then preserved with 70% ethanol. Prepared samples were homogenized using a Nissei ACE homogenizer (Nihonseiki Kaisha LTD, Tokyo, Japan) for 5 min until the tissue was well-mixed. Subsamples ($2.5 \mu\text{l}$) of the coral tissue homogenate were immediately placed on each chamber of a Neubauer-improved bright line hemocytometer (Marien Feld, Germany) to count the number of zooxanthellae under 40x magnification. Four corner squares of each chamber were counted as one replication. To calculate the mean zooxanthellae density per cm^2 , the average total of zooxanthellae cell counted on each square was multiplied by 10^4 ml (the volume of each chamber) and




Figure 1 Study site and sampling period. Ko Tao Mo, Sattahip, upper Gulf of Thailand. The red arrows showed the sampling times.

Full-size [DOI: 10.7717/peerj.13114/fig-1](https://doi.org/10.7717/peerj.13114/fig-1)

the dilution then divided by the coral tissue area (cm²). The average zooxanthellae density of two coral species from three different seasons were tested using a two-way ANOVA and Post Hoc tests.

DNA extraction and PCR amplification

Small pieces of coral preserved in 95% ethanol were dried and ground using a pestle and mortar to create a coarse powder. Each powdered coral was used to extract the genomic DNA of zooxanthellae using a DNeasy Plant Mini Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions with minor modification). Crude DNA was analyzed using a NanoDrop 2000c (Thermo Fisher Scientific, Waltham, MA, USA) to examine the quality and concentration of DNA. Extracted DNA was used as a PCR template to amplify the DNA of the Symbiodiniaceae. The PCR amplifications were performed using primers for the internal transcribed spacers 1 and 2 (ITS-1 and ITS-2) genes. The targeted ITS-1 and ITS-2 regions were amplified using the specific primers for Symbiodiniaceae species identification including forward: r18Sf (5'-CGCTCTCCGATCTCTGGAAAGTTTCATGAACCTTAT 3'-) and reverse: Sym28Sr-1st (5'-TGCTCTCCGATCTGACCTTGTRTGACTTCATGCTA -3'). PCR was carried out using AllInOneCycler™ (BIONEER, Inc., Oakland, CA, USA) with a reaction volume of 10 µl mixture for each sample which consisted of 0.2 µl PrimeSTAR GXL DNA Polymerase, 2 µl Taq buffer, 0.8 µl dNTP of Takara (TAKARA Bio Inc., Shiga, Japan), 1 µl of each forward and reverse primer, 4 µl of 0.1 µm PCR

grade water (Sigma[®] Life Science) and 1 μl of DNA template. The PCR conditions were 3 min at 94 °C, followed by 33 cycles of 98 °C for 10 s, 50 °C for 0.15 s, 68 for 60 s, and a final extension step of 3 m at 68 °C. PCR products were cleaned using the FavorPrep[®] gel/PCR Purification Kit (FAVORGEN Biotech Corp., Wembley, WA, Australia) following the manufacturer's protocol. Each purified sample was then used for the second PCR. The unique short nucleotides barcode (Eurofins Scientific, Luxembourg, Germany) including forward and reverse were performed. Here 90 pairs of different primers (90 coral samples) were used to include in the second PCR mixture. Each 10 μl PCR sample included the following components: 0.24 μl PrimeSTAR GXL DNA Polymerase, 2.4 μl Taq buffer, 0.96 μl dNTP of Takara (TAKARA Bio Inc., Shiga, Japan), 1 μl (0.1 μM) of each unique primer combination of forward and reverse primers, 3.4 μl Sigma water and 1 μl of purified PCR product. The PCR conditions were 3 min at 94 °C, followed by 13 cycles of 98 °C 10 s, 68 °C 60 s. Second PCR products were run in a 1.5% agarose gel with EcoDye Nucleic Acid Staining Solution (BIOFACT Co., Ltd., Daejeon, Korea) to check the targeted size of DNA.

Next-generation sequencing and data analysis

Each successful PCR amplicon was pooled together in an equimolar concentration for sequencing. Pooled samples were cleaned using SPRI Based Size Selection (Beckman Coulter, Inc, USA). DNA libraries then were quantified on Agilent Bioanalyzer 2100 (Agilent Technologies, Palo Alto, CA, United States) and qPCR (Thermo Fisher Scientific, Waltham, MA, USA) to examine the concentration and average DNA fragment size (bp). We used 4nM of DNA libraries and made a dilution to 8 pM followed by sequencing on an Illumina Mi-Seq platform (Illumina, San Diego, CA, United States) by the Atmosphere and Ocean Research Institute, The University of Tokyo, Japan. The analysis pipeline and analytical procedures were carried out following [Shinzato et al. \(2018\)](#). The raw data were submitted to the NCBI Sequence Read Archive (BioProject ID: [PRJNA747703](#)). For the statistical analysis of difference in Symbiodiniaceae diversity, Shannon and Simpson indices were calculated to assess the level of alpha diversity in both corals ([Simpson, 1949](#)). The Kruskal–Wallis test was performed to test the difference in the Symbiodiniaceae community during different seasons. A permutation multivariate analysis of variance (PERMANOVA) test was used to compare the Symbiodiniaceae community compositions among coral species. To visualize the beta diversity of Symbiodiniaceae community structure among coral hosts, non-metric multidimension scaling (nMDS) based on Bray–Curtis dissimilarity was plotted using Phyloseq and VEGAN packages in R software ([McMurdie & Holmes, 2013](#); [Oksanen et al., 2020](#)).

RESULTS

Local environment condition

The local conditions at Ko Tao Mo reef were continuously recorded during each season over a year. The seasonal minimum and maximum temperatures were recorded in January (26.90 °C) and May (32.18 °C) contrasting with light intensity, which was lowest in August and September (1 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and highest in April (342 $\mu\text{mol m}^{-2} \text{s}^{-1}$), respectively. An average local light intensity and temperature in each season was found to




Figure 2 Average light intensity and temperature from winter to rainy seasons at Ko Tao Mo, Chon Buri, Upper Gulf of Thailand during November 2017 to October 2018.

Full-size  DOI: [10.7717/peerj.13114/fig-2](https://doi.org/10.7717/peerj.13114/fig-2)

be 81.25 ± 2.50 , 121.86 ± 8.49 , $55.90 \pm 3.84 \mu\text{mol m}^{-2} \text{s}^{-1}$ and 28.80 ± 0.11 , 31.06 ± 0.06 and $30.67 \pm 0.06 \text{ }^\circ\text{C}$ following winter (November–February), summer (March–June) and the rainy season (July–October), respectively (Fig. 2). While both light intensity and temperature were highest in the summer, field observations found no sign of coral bleaching (though a slightly paler color was recorded in both *A. humilis* and *P. cf. damicornis* colonies) during 2018.

Zooxanthellae density

Average zooxanthellae density (ZDs) in *A. humilis* and *P. cf. damicornis* across the three seasons ranged from 0.66×10^6 to 3.69×10^6 cells/cm². The mean ZDs in *P. cf. damicornis* were twice as high ($3.45 \pm 0.62 \times 10^6$ cells/cm²) as *A. humilis* ($1.75 \pm 0.42 \times 10^6$ cells/cm²) (Supplementary 2). The variation of physical factors in different seasons was found to significantly impact ZDs especially in *A. humilis*. Pearson correlation among zooxanthellae cell density, light intensity and temperature in this study is presented in Table 1. The density of zooxanthellae cells showed strong negative correlations with light intensity and a weak inverse correlation between temperature. The mean ZDs in both species showed the lowest number during the summer with *A. humilis* having $0.8 \pm 0.18 \times 10^6$ cells/cm² and *P. cf. damicornis* having $3.66 \pm 0.42 \times 10^6$ cells/cm². The highest ZDs in *P. cf. damicornis* were found to occur in winter (3.69×10^6 cells/cm²) while the highest in *A. humilis* was in the rainy season ($2.21 \pm 0.03 \times 10^6$ cells/cm²) (Fig. 3). However, the comparison of ZDs between winter and rainy season in both corals was not found to be significantly different ($p > 0.05$) (Supplementary 3).

Table 1 Summary of two-way ANOVAs analyses on Symbiodiniaceae density in *A. humilis* and *P. cf. damicornis* at different local light intensities and temperature.

	Light intensity ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	Temperature ($^{\circ}\text{C}$)	<i>A. humilis</i> ($\times 10^6$ cells/cm 2)	<i>P. damicornis</i> ($\times 10^6$ cells/cm 2)
Light intensity	1			
Temperature	.610	1		
ZDs of <i>A. humilis</i> ($\times 10^6$ cells/cm 2)	-0.98**	-0.64	1	
ZDs of <i>P. damicornis</i> ($\times 10^6$ cells/cm 2)	-0.76*	-0.53	0.71	1

Notes.

** Correlation is significant at the 0.01 level.

* Correlation is significant at the 0.05 level.




Figure 3 Box plots describing average of zooxanthellae densities of *A. humilis* and *P. cf. damicornis* during the winter, summer and rainy seasons. The median abundances and the interquartile ranges have been indicated in the plots.

Full-size DOI: 10.7717/peerj.13114/fig-3

Diversity and composition of Symbiodiniaceae of *A. humilis* and *P. cf. damicornis* corals

Genomic DNA was obtained from 80 colonies of two coral species from three seasons. Sequencing of the ITS-1 and ITS-2 amplicons yielded 3,450,698 raw demultiplexed reads. Adaptor and quality trimmed reads resulted in 1,488,632 raw contigs and yielded 1,137,320 contigs after chimera removed reads from 80 individuals. An average 14,217 contigs/sample (Supplementary 4). Based on the GeoSymbio database, two genera of Symbiodiniaceae were detected: *Cladocopium* (14 species) and *Durusdinium* (six species). The alpha diversity indices (Shannon and Simpson) of coral showed that *A. humilis* appeared to associate with Symbiodiniaceae communities with higher levels of diversity than *P. cf. damicornis* (Fig. 4) (Supplementary 4). A total of 14 species were identified in the two coral species, with *A. humilis* consisting of two dominant species including Cspc_C3 (56.39%), C3w (33.62%) and other minor species C93type1 (4.42%), Cspf (3.59%) D1 (1.03%) and other species




Figure 4 Box plots showing alpha diversity indices of Symbiodiniaceae community from two coral species, *Acropora humilis* ($n = 15$) and *Pocillopora cf. damicornis* ($n = 15$) in different seasons. Shannon and Simpson indexes reflects the evenness and diversity of Symbiodiniaceae species, respectively. Median values and interquartile ranges were indicated in the plots. Filled circles represent outliers. Differences were assessed by non-parametric Kruskal–Wallis test, and p -values are shown in each box ($p > 0.05$).

Full-size DOI: 10.7717/peerj.13114/fig-4

(0.95%). On the other hand, the majority of *Durusdinium* was found in *P. cf. damicornis* following D1 (95.58%), D6 (1.01%) D10 (2.70%) and other species (0.71%) (Fig. 5). A Bray-Curtis based nMDS plot illustrated a significant difference in Symbiodiniaceae communities between *A. humilis* and *P. cf. damicornis* (PERMANOVA, $p < 0.001$). The number of valid alignments of Symbiodiniaceae species across the different seasons remained relatively consistent in *A. humilis* and *P. cf. damicornis* throughout the year but a greater dominance of *Durusdinium* D1 was found in one colony of *A. humilis* during the rainy season (Fig. 6).

DISCUSSION

Our study confirmed that each coral genus contained significantly different numbers of zooxanthellae. However, the average zooxanthellae densities in both *A. humilis* and *P. cf. damicornis* (1.75 and 3.44×10^6 cells/cm²) were within the range shown in other studies in the Pacific and Indian ocean with approximately $1\text{--}3 \times 10^6$ cells/cm² (Fitt et al., 2001; Li et al., 2008; Pillay, Willis & Terashima, 2005; Qin et al., 2019; Stimson, 1997; Xu et al., 2017; Zhou et al., 2017). Several factors can apparently determine the number of zooxanthellae cells in each coral species (Fitt et al., 2009; Yuyama et al., 2016). One such characteristic is tissue thickness, which is correlated with the capacity to absorb light (A_{\max}) by each coral (Scheufen, Iglesias-Prieto & Enríquez, 2017). For example, thinner tissue layers and smaller polyps such as are found in *Acropora* contained lower numbers




Figure 5 Relative abundance of Symbiodiniaceae communities in *A. humilis* and *P. cf. damicornis* across different seasons in 2018.

Full-size [DOI: 10.7717/peerj.13114/fig-5](https://doi.org/10.7717/peerj.13114/fig-5)




Figure 6 Non-metric multidimensional scaling (nMDS) plotting of the Symbiodiniaceae species composition of two coral species based on of Bray-Curtis dissimilarity indices.

Full-size [DOI: 10.7717/peerj.13114/fig-6](https://doi.org/10.7717/peerj.13114/fig-6)

of zooxanthellae than corals from genera such as *Porites* and *Stylophora* which has a thicker tissue, such that the thicker tissue corals require a higher symbiont densities or pigmentation to reach A_{max} (Ben-Zvi, Eyal & Loya, 2015; Li et al., 2008; Loya et al., 2001; Scheufen, Iglesias-Prieto & Enríquez, 2017; Tambutté et al., 2007). Moreover, corals harbored different species of Symbiodiniaceae, which perform differently in term of photosynthetic activity (Cantin et al., 2009; Yuyama et al., 2018). Generally, *Durudinium* has a relatively lower potential to provide photosynthetically fixed carbon to the coral host compared with *Cladocopium* (Sproles et al., 2020). Other studies have confirmed that the same coral species which contained different genera of Symbiodiniaceae show a lower concentration of photopigments per zooxanthellae cell in *Durudinium*-colonies, resulting in a higher number of zooxanthellae cells required by those corals (Wall et al., 2020; Yuyama

& Higuchi, 2014). This may explain the different symbiont densities detected in *A. humilis* (thin tissue) and *P. cf. damicornis* (thick tissue) in this study. It is however important to state that the variation of zooxanthellae density also depends on sampling depth, coral species, local environmental factors (Al-Hammady, 2013; Brown et al., 1999b; Ladrière et al., 2014; Stimson, 1997; Xu et al., 2017), zooxanthellae count preparation method (McCowan et al., 2011) and interval time of sample collections.

Seasonal changes have been shown to influence zooxanthellae density variation in both corals (*A. humilis* and *P. cf. damicornis*), especially because the high light intensity and temperature during summer are negatively correlated with zooxanthellae densities (Hill & Ralph, 2007; Pillay, Willis & Terashima, 2005). Rising temperature and light intensity were found to cause a reduction in zooxanthellae cell counts in several shallow water corals in Thailand (Andaman Sea) (Brown et al., 1999b). Likewise, assessments of zooxanthellae function have also been carried out *ex-situ* based on colonies found near the present study site of Ko Tao Mo. The corals, *A. millepora* and *Platygyra sinensis*, were shown to have a statistically significant reduction in photosynthetic rate (F_v/F_m) under higher light intensities (Kuanui et al., 2020; Kuanui et al., 2015). Thermal stress during coral exposure to high UV radiation and elevated temperature resulted in an overproduction of reactive oxygen species (ROS) by intracellular endosymbiont cells (Nielsen, Petrou & Gates, 2018; Vidal-Dupiol et al., 2009). The enhanced production of ROS and antioxidant expression in corals not only causes irreversible DNA damage but also has the consequence of increased metabolic activity, which becomes toxic to the host (Downs et al., 2002; Wang et al., 2017). Therefore, scleractinian corals respond to their physiological stress by expelling the endosymbiont cells to minimize intracellular damage and balancing of energy dissipation efficiency during environmental stress (Kemp et al., 2014; Nielsen, Petrou & Gates, 2018; Weis, 2008). On the other hand, corals established higher zooxanthellae density during winter in some subtropical and tropical reef regions (Costa, Sassi & Amaral, 2004; Ferrier-Pagès et al., 2011; Fitt et al., 2000; Pillay, Willis & Terashima, 2005; Stimson, 1997), which may be linked to an optimum range of light and temperature for photosynthetic activity in coral at this particular site (Kleypas, McManus & MeÑez, 1999; Schoepf et al., 2015). Pillay, Willis & Terashima (2005) also reported that the higher zooxanthellae densities were recorded during winter, which may be due to the physiological seasonal adaptations of corals during low light conditions for harvesting capacity.

The Symbiodiniaceae community in the present study was similar to previous studies in Thailand, with only two genera, *Cladocopium* and *Durusdinium*, being found both in the Gulf of Thailand and Andaman Sea but different coral genera were identified (Chankong et al., 2020; Pootakham et al., 2018). The community structure of Symbiodiniaceae with particular coral species also showed a similar pattern to those in Xinyi and Palau reefs, in which, *Acropora* spp. hosted *Cladocopium* while *Pocillopora* spp. contained *Durusdinium* D1 almost exclusively (Qin et al., 2019; Wham, Ning & LaJeunesse, 2017). Nevertheless, the different patterns of the same coral genus associated with Symbiodiniaceae genus/species is known to occur across both local scales and wide geographic zones (Rodriguez-Lanetty et al., 2001; Ulstrup & Van Oppen, 2003; Van Oppen et al., 2005). Several studies in the Great Barrier Reef (Lizard Island) were found to show the opposite

trend of Symbiodiniaceae community in Acroporidae and Pocilloporidae, with *Acropora* being associated several species of Symbiodiniaceae, such as *Symbiodinium* (A3_302), *Cladocopium* (C1_769, C42.2_1367, and C3nn_1031) while *Pocillopora* associated with almost 100% of *Cladocopium* (C1_769) (Ziegler et al., 2018).

This study indicates that corals showed a strong specificity to a particular Symbiodiniaceae genus which may correlate with the pattern of symbiont acquisition (Fabina et al., 2012; LaJeunesse et al., 2004; Stat et al., 2008). The coral-Symbiodiniaceae association may be influenced by several factor including environmental variability, physiology, life history, and reproductive background (LaJeunesse et al., 2004; LaJeunesse et al., 2010). Approximately 80–90% of reef building corals in the Indo-Pacific region are broadcast spawners, which produce zooxanthellae-free gametes (Richmond & Hunter, 1990; Yuyama et al., 2016). Generally, *Acropora* spp. are considered to horizontally transmit symbionts and can harbor several species of Symbiodiniaceae to cope with environmental stress, such as was found for *A. humilis* in this study (Gong et al., 2019; LaJeunesse et al., 2010; Putnam et al., 2012; Quigley, Willis & Bay, 2017). The offspring of these hosts probably reflected local environmental conditions during the early life stages when the symbiosis with Symbiodiniaceae was initially developed (LaJeunesse et al., 2004). Free-living Symbiodiniaceae cells, both in the water column and sediment, are an important natural pool for corals during larval or primary polyp stage and/or under critical conditions (Adams, Cumbo & Takabayashi, 2009; Hoegh-Guldberg et al., 2008). Work by Yamashita et al. (2014) found that *Cladocopium* were detected mostly in the sediment while *Symbiodinium* and *Durusdinium* were found in the water column. This could explain how Symbiodiniaceae cells in the sediment play a significant role in symbiont uptake in *A. humilis*. On the other hand, *Pocillopora* has been shown to associate with *Durusdinium*. *Pocillopora* cf. *damicornis* in Thailand has been reported as a brooding coral, which releases planulae containing Symbiodiniaceae cells (Kuanui et al., 2008). The Symbiodiniaceae community of these corals' offspring are hypothesized to be more similar to their mother colony due to the vertical symbiont transmission process (Hirose & Hidaka, 2006; Quigley et al., 2018). Consequently, most VT corals including *P.* cf. *damicornis* possess greater stability with Symbiodiniaceae species across biogeographic gradients including Thai waters with almost 100% association with *Durusdinium* D1 (Chankong et al., 2020; Muller-Parker & Davy, 2001; Pootakham et al., 2021; Thornhill, Fitt & Schmidt, 2006). However, there is evidence that some brooding corals exhibit a mixed-mode acquisition of Symbiodiniaceae (vertical transmission and horizontal symbiont acquisition). Consequently, these corals may become more tolerant to stressful conditions than previously thought by up-taking the resilient Symbiodiniaceae species from the environment (Epstein et al., 2019; Quigley et al., 2018; Quigley, Willis & Kenkel, 2019).

Acropora humilis in this trial hosted *Cladocopium* species and were found to be more susceptible to bleaching when higher light intensity was presented. Similarly, several studies have confirmed that *Acropora* spp. tend to be more susceptible to bleaching, which may be linked to the species of Symbiodiniaceae with which it associated (Hoogenboom et al., 2017; Putchim et al., 2017; Sakai, Singh & Iguchi, 2019), whereas *P.* cf. *damicornis* containing *Durusdinium* showed fewer effects on the corals in the different seasons.

Distinct genera/species of Symbiodiniaceae have been shown to perform differently under changing environments. For example, *Durusdinium* and *Symbiodinium* are considered to be stress-tolerant and/or opportunistic species (ie., warm, high turbidity, high irradiance) (LaJeunesse, 2002; Levas et al., 2013; Rowan, 2004; Wham, Ning & LaJeunesse, 2017) whereas corals associated with *Cladocopium* species exhibited more susceptibility during stressful conditions in some coral reef regions (Suwa, Hirose & Hidaka, 2008; Thinesh et al., 2019).

Our results revealed that the seasonal variation does not affect the dominant genus/species of Symbiodiniaceae in both *A. humilis* and *P. cf. damicornis* corals. Most corals have not been shown to change their species or community composition of Symbiodiniaceae, despite continued changes in environmental conditions (Coffroth et al., 2010; Goulet, 2006; Keshavmurthy et al., 2020; Lewis & Coffroth, 2004). There are several studies that have shown a similar trend along tropical to temperate regions (Berkelmans & Van Oppen, 2006; Cunning, Silverstein & Baker, 2018; Rouze et al., 2017). Ng & Ang (2016) who investigated labeled colonies of *Platygyra acuta* and *Porites* spp. from Hong Kong reefs in different seasons found that there was no change in the Symbiodiniaceae species in all corals. Similarly, *Porites lutea* in the Andaman Sea, Thailand, retain a stable Symbiodiniaceae community structure (C116, C15f, C15g, C15 h, C15.2) even in bleached corals (Pootakham et al., 2018). Although, Kagoshima Bay, Japan has a high variability of temperature, there was no variation of Symbiodiniaceae (C1, C3-related) in *Zoanthus saensibaricus* (zoanthids) across twelve months of the experiment (Reimer, 2007). In this way, Symbiodiniaceae shuffling or switching in corals might not be a common mechanism to survive under environmental stress (Goulet, 2006; Keshavmurthy et al., 2020).

Several possible hypotheses could explain why corals maintain a stable community of Symbiodiniaceae during environmental stress; 1) avoiding intraspecific competition among Symbiodiniaceae species that could lead to disequilibrium of nutrients within the coral host and affect coral growth, 2) coral response to historical environment stressors may promote stable Symbiodiniaceae associations, 3) higher composition of some Symbiodiniaceae species may effect coral-cell metabolism, 4) there may be no severe effect on corals after experiencing stressful conditions (Jones & Berkelmans, 2011; Klepac et al., 2015; LaJeunesse et al., 2010; Putnam et al., 2012) or 5) limitation of high-stress tolerant Symbiodiniaceae genotypes within the host coral and/or environment. However, the variation of Symbiodiniaceae diversity and/or communities may be driven by several factors including local environmental variability, coral species background, Symbiodiniaceae supply sources such as seawater, sediment etc. and investigation techniques (Baird et al., 2009; LaJeunesse et al., 2010; Ng & Ang, 2016; Sweet, 2014; Thornhill et al., 2014; Van Oppen et al., 2005). Although the corals-Symbiodiniaceae associations in the upper Gulf of Thailand were found to be stable during the studied year, we predict that increasing severe coral bleaching in the future may result in acclimatization or adaptation of some corals by Symbiodiniaceae shuffling within hosts.

CONCLUSIONS

Both *A. humilis* and *P. cf. damicornis* in this study exhibited the lowest zooxanthellae density during the summer, which is likely caused by light intensity. In addition, the corals had a strong relationship with particular Symbiodiniaceae species, specific to each species, and showed no variation of Symbiodiniaceae species throughout the year. The corals in the upper Gulf of Thailand may select the appropriate genus/species of Symbiodiniaceae in response to local environmental stressors. The results of this study provided some information on the coral-Symbiodiniaceae relationship between seasons, which may be applied to predicting the potential adaptation of corals in localized reef environments. However, more coral species and study sites are needed to further study the seasonal variation on coral-Symbiodiniaceae in Thailand.

ACKNOWLEDGEMENTS

We would like to thank the Plant Genetic Conservation Project under the Royal Initiative of Her Royal Highness Princess Maha Chakri Sirindhorn and the Naval Special Warfare Command, Royal Thai Fleet, Royal Thai Navy for their assisting in the fieldwork. We also acknowledged the UNESCO/IOC-WESTPAC and UNESCO Japanese Funds in Trust for assisting in the meetings related to the coral conservation project.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research was supported by the 90th Anniversary of Chulalongkorn University Fund (Ratchadaphiseksomphot Endowment Fund), the Royal Golden Jubilee Ph.D. program (Grant No. PHD/0126/2558), the Thailand Research Fund, the NRCT-JSPS Core to Core Program, CREPSUM JPJSCCB20200009, the Thailand Science Research and Innovation Fund Chulalongkorn University (CU_FRB65_dis (3)_091_23_23)), and Mubadala Petroleum (Thailand) Limited. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

The 90th Anniversary of Chulalongkorn University Fund (Ratchadaphiseksomphot Endowment Fund).

The Royal Golden Jubilee Ph.D. program: PHD/0126/2558.

Thailand Research Fund.

NRCT-JSPS Core to Core Program.

CREPSUM JPJSCCB20200009.

Thailand Science Research and Innovation Fund Chulalongkorn University:

CU_FRB65_dis (3)_091_23_23.

Mubadala Petroleum (Thailand) Limited.

Competing Interests

Suchana Chavanich is an Academic Editor for PeerJ.

Author Contributions

- Suppakarn Jandang conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Voranop Viyakarn conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Yuki Yoshioka performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Chuya Shinzato performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Suchana Chavanich conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The Symbiodiniaceae sequences are available at GenBank: [PRJNA747703](https://www.ncbi.nlm.nih.gov/GenBank/PRJNA747703).

Data Availability

The following information was supplied regarding data availability:

The number of sequencing reads and alpha diversity indices of two coral species, *Acropora humilis* and *Pocillopora damicornis*, is available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13114#supplemental-information>.

REFERENCES

- Adams L, Cumbo V, Takabayashi M. 2009.** Exposure to sediment enhances primary acquisition of *Symbiodinium* by asymbiotic coral larvae. *Marine Ecology Progress Series* 377:149–156 DOI 10.3354/meps07834.
- Al-Hammady MAM. 2013.** The effect of zooxanthellae availability on the rates of skeletal growth in the Red Sea coral *Acropora hemprichii*. *The Egyptian Journal of Aquatic Research* 39(3):177–183 DOI 10.1016/j.ejar.2013.10.005.
- Baird AH, Bhagooli R, Ralph PJ, Takahashi S. 2009.** Coral bleaching: the role of the host. *Trends in Ecology & Evolution* 24(1):16–20 DOI 10.1016/j.tree.2008.09.005.
- Baker AC. 2003.** Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and biogeography of *Symbiodinium*. *Annual Review of Ecology, Evolution, and Systematics* 34(1):661–689 DOI 10.1146/annurev.ecolsys.34.011802.132417.

- Baker DM, Weigt L, Fogel M, Knowlton N. 2013.** Ancient DNA from coral-hosted *Symbiodinium* reveal a static mutualism over the last 172 years. *PLOS ONE* **8**(2):e55057 DOI [10.1371/journal.pone.0055057](https://doi.org/10.1371/journal.pone.0055057).
- Baskett ML, Gaines SD, Nisbet RM. 2009.** Symbiont diversity may help coral reefs survive moderate climate change. *Ecological Applications* **19**(1):3–17 DOI [10.1890/08-0139.1](https://doi.org/10.1890/08-0139.1).
- Bednarz V, Cardini U, Van Hoytema N, Al-Rshaidat M, Wild C. 2015.** Seasonal variation in dinitrogen fixation and oxygen fluxes associated with two dominant zooxanthellate soft corals from the northern Red Sea. *Marine Ecology Progress Series* **519**:141–152 DOI [10.3354/meps11091](https://doi.org/10.3354/meps11091).
- Ben-Zvi O, Eyal G, Loya Y. 2015.** Light-dependent fluorescence in the coral *Galaxea fascicularis*. *Hydrobiologia* **759**(1):15–26 DOI [10.1007/s10750-014-2063-6](https://doi.org/10.1007/s10750-014-2063-6).
- Berkelmans R, Van Oppen MJ. 2006.** The role of zooxanthellae in the thermal tolerance of corals: a ‘nugget of hope’ for coral reefs in an era of climate change. *Proceedings Biological Sciences* **273**(1599):2305–2312 DOI [10.1098/rspb.2006.3567](https://doi.org/10.1098/rspb.2006.3567).
- Blackall LL, Wilson B, Van Oppen MJH. 2015.** Coral—the world’s most diverse symbiotic ecosystem. *Molecular Ecology* **24**(21):5330–5347 DOI [10.1111/mec.13400](https://doi.org/10.1111/mec.13400).
- Boillard A, Dubé CE, Gruet C, Mercière A, Hernandez-Agreda A, Derome N. 2020.** Defining coral bleaching as a microbial dysbiosis within the coral holobiont. *Microorganisms* **8**(11):1682 DOI [10.3390/microorganisms8111682](https://doi.org/10.3390/microorganisms8111682).
- Brown BE, Ambarsari I, Warner ME, Fitt WK, Dunne R, Gibb SW, Cummings DG. 1999a.** Diurnal changes in photochemical efficiency and xanthophyll concentrations in shallow water reef corals: evidence for photoinhibition and photoprotection. *Coral Reefs* **18**:99–105 DOI [10.1007/s003380050163](https://doi.org/10.1007/s003380050163).
- Brown BE, Ambarsari I, Warner ME, Fitt WK, Dunne RP, Gibb SW, Cummings DG. 1999b.** Diurnal changes in photochemical efficiency and xanthophyll concentrations in shallow water reef corals: evidence for photoinhibition and photoprotection. *Coral Reefs* **18**(2):99–105 DOI [10.1007/s003380050163](https://doi.org/10.1007/s003380050163).
- Cantin NE, Van Oppen MJH, Willis BL, Mieog JC, Negri AP. 2009.** Juvenile corals can acquire more carbon from high-performance algal symbionts. *Coral Reefs* **28**(2):405–414 DOI [10.1007/s00338-009-0478-8](https://doi.org/10.1007/s00338-009-0478-8).
- Chankong A, Kongjandtre N, Senanan W, Manthachitra V. 2020.** Community composition of Symbiodiniaceae among four scleractinian corals in the eastern Gulf of Thailand. *Regional Studies in Marine Science* **33**:100918 DOI [10.1016/j.rsma.2019.100918](https://doi.org/10.1016/j.rsma.2019.100918).
- Chen B, Yu K, Qin Z, Liang J, Wang G, Huang X, Wu Q, Jiang L. 2020.** Dispersal, genetic variation, and symbiont interaction network of heat-tolerant endosymbiont *Durusdinium trenchii*: insights into the adaptive potential of coral to climate change. *Science of the Total Environment* **723**:138026 DOI [10.1016/j.scitotenv.2020.138026](https://doi.org/10.1016/j.scitotenv.2020.138026).
- Coffroth MA, Poland DM, Petrou EL, Brazeau DA, Holmberg JC. 2010.** Environmental symbiont acquisition may not be the solution to warming seas for reef-building corals. *PLOS ONE* **5**(10):e13258 DOI [10.1371/journal.pone.0013258](https://doi.org/10.1371/journal.pone.0013258).
- Costa CF, Sassi R, Amaral F. 2004.** Population density and photosynthetic pigment content in symbiotic dinoflagellates in the Brazilian scleractinian coral *Montastrea*

cavernosa. *Brazilian Journal of Oceanography* 52:93–99
DOI 10.1590/S1679-87592004000200001.

- Császár NBM, Ralph PJ, Frankham R, Berkelmans R, Van Oppen MJH. 2010. Estimating the potential for adaptation of corals to climate warming. *PLOS ONE* 5(3):e9751–e9751 DOI 10.1371/journal.pone.0009751.
- Cunning R, Silverstein R, Baker A. 2015. Investigating the causes and consequences of symbiont shuffling in a multi-partner reef coral symbiosis under environmental change. *Proceedings. Biological Sciences* 282:1–8 DOI 10.1098/rspb.2014.1725.
- Cunning R, Silverstein R, Baker A. 2018. Symbiont shuffling linked to differential photochemical dynamics of *Symbiodinium* in three Caribbean reef corals. *Coral Reefs* 37:145–152 DOI 10.1007/s00338-017-1640-3.
- Davies S, Ries J, Marchetti A, Castillo K. 2018. Symbiodinium functional diversity in the coral *Siderastrea siderea* is influenced by thermal stress and reef environment, but not ocean acidification. *Frontiers in Marine Science* 5:150 DOI 10.3389/fmars.2018.00150.
- DeSalvo MK, Sunagawa S, Fisher PL, Voolstra CR, Iglesias-Prieto R, Medina M. 2010. Coral host transcriptomic states are correlated with *Symbiodinium* genotypes. *Molecular Ecology* 19(6):1174–1186 DOI 10.1111/j.1365-294X.2010.04534.x.
- Dilworth J, Caruso C, Kahkejian V, Baker A, Drury C. 2021. Host genotype and stable differences in algal symbiont communities explain patterns of thermal stress response of *Montipora capitata* following thermal pre-exposure and across multiple bleaching events. *Coral Reefs* 40:1–13 DOI 10.1007/s00338-020-02024-3.
- Downs C, Fauth J, Halas J, Dustan P, Bemiss J, Woodley C. 2002. Oxidative stress and seasonal coral bleaching. *Free Radical Biology & Medicine* 33:533–543 DOI 10.1016/S0891-5849(02)00907-3.
- Eckert RJ, Reaume AM, Sturm AB, Studivan MS, Voss JD. 2020. Depth influences Symbiodiniaceae associations among *Montastraea cavernosa* corals on the Belize Barrier Reef. *Frontiers in Microbiology* 11:1–13 DOI 10.3389/fmicb.2020.00518.
- Epstein HE, Torda G, Munday PL, Van Oppen MJH. 2019. Parental and early life stage environments drive establishment of bacterial and dinoflagellate communities in a common coral. *The ISME Journal* 13(6):1635–1638 DOI 10.1038/s41396-019-0358-3.
- Fabina NS, Putnam HM, Franklin EC, Stat M, Gates RD. 2012. Transmission mode predicts specificity and interaction patterns in coral-*Symbiodinium* networks. *PLOS ONE* 7(9):e44970 DOI 10.1371/journal.pone.0044970.
- Fabricius K, Mieog J, Colin P, Idip D, Van Oppen M. 2004. Identity and diversity of coral endosymbionts (zooxanthellae) from three Palauan reefs with contrasting bleaching, temperature and shading histories. *Molecular Ecology* 13:2445–2458 DOI 10.1111/j.1365-294X.2004.02230.x.
- Ferrier-Pagès C, Gattuso J-P, Cauwet G, Jaubert J, Allemand D. 1998. Release of dissolved organic matter and nitrogen by the zooxanthellate coral *Galaxa fascicularis*. *Marine Ecology Progress Series* 172:263–274.
- Ferrier-Pagès C, Peirano A, Abbate M, Cocito S, Negri A, Rottier C, Riera P, Rodolfo-Metalpa R, Reynaud S. 2011. Summer autotrophy and winter heterotrophy in

- the temperate symbiotic coral *Cladocora caespitosa*. *Limnology and Oceanography* **56**:1429–1438 DOI [10.4319/lo.2011.56.4.1429](https://doi.org/10.4319/lo.2011.56.4.1429).
- Fisher PL, Malme MK, Dove S. 2012.** The effect of temperature stress on coral–*Symbiodinium* associations containing distinct symbiont types. *Coral Reefs* **31**(2):473–485 DOI [10.1007/s00338-011-0853-0](https://doi.org/10.1007/s00338-011-0853-0).
- Fitt W, Brown B, Warner M, Dunne R. 2001.** Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs* **20**:51–65 DOI [10.1007/s003380100146](https://doi.org/10.1007/s003380100146).
- Fitt W, Gates R, Hoegh-Guldberg O, Bythell J, Jatkar A, Grottoli A, Gomez Cabrera KL, Fisher P, Lajuenesse T, Pantos O, Iglesias-Prieto R, Franklin D, Rodrigues L, Torregiani J, Van Woesik R, Lesser M. 2009.** Response of two species of Indo-Pacific corals, *Porites cylindrica* and *Stylophora pistillata*, to short-term thermal stress: the host does matter in determining the tolerance of corals to bleaching. *Journal of Experimental Marine Biology and Ecology* **373**:102–110 DOI [10.1016/j.jembe.2009.03.011](https://doi.org/10.1016/j.jembe.2009.03.011).
- Fitt WK, McFarland FK, Warner ME, Chilcoat GC. 2000.** Seasonal patterns of tissue biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral bleaching. *Limnology and Oceanography* **45**(3):677–685 DOI [10.4319/lo.2000.45.3.0677](https://doi.org/10.4319/lo.2000.45.3.0677).
- Ghavam Mostafavi P, Fatemi SMR, Shahhosseiny MH, Hoegh-Guldberg O, Loh WKW. 2007.** Predominance of clade D *Symbiodinium* in shallow-water reef-building corals off Kish and Larak Islands (Persian Gulf, Iran). *Marine Biology* **153**(1):25–34 DOI [10.1007/s00227-007-0796-8](https://doi.org/10.1007/s00227-007-0796-8).
- Glynn P. 1993.** Glynn, P. W. Coral bleaching: ecological perspective. *Coral Reefs* **12**:1–17 DOI [10.1007/BF00303779](https://doi.org/10.1007/BF00303779).
- Gong S, Xu L, Yu K, Zhang F, Li Z. 2019.** Differences in Symbiodiniaceae communities and photosynthesis following thermal bleaching of massive corals in the northern part of the South China Sea. *Marine Pollution Bulletin* **144**:196–204 DOI [10.1016/j.marpolbul.2019.04.069](https://doi.org/10.1016/j.marpolbul.2019.04.069).
- Goulet T. 2006.** Most corals may not change their symbionts. *Marine Ecology Progress Series* **321**:1–7 DOI [10.3354/meps321001](https://doi.org/10.3354/meps321001).
- Hill R, Ralph PJ. 2007.** Post-bleaching viability of expelled zooxanthellae from the scleractinian coral *Pocillopora damicornis*. *Marine Ecology Progress Series* **352**:137–144 DOI [10.3354/meps07159](https://doi.org/10.3354/meps07159).
- Hinrichs S, Patten NL, Allcock RJN, Saunders SM, Strickland D, Waite AM. 2013.** Seasonal variations in energy levels and metabolic processes of two dominant *Acropora* species (*A. spicifera* and *A. digitifera*) at Ningaloo Reef. *Coral Reefs* **32**(3):623–635 DOI [10.1007/s00338-013-1027-z](https://doi.org/10.1007/s00338-013-1027-z).
- Hirose M, Hidaka M. 2006.** Early development of zooxanthella-containing eggs of the corals *Porites cylindrica* and *Montipora digitata*: the endodermal localization of zooxanthellae. *Zoological Science* **23**(10):873–881 DOI [10.2108/zsj.23.873](https://doi.org/10.2108/zsj.23.873).

- Hoegh-Guldberg O, Mumby P, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell C, Sale P, Edwards A, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury R, Dubi A, Hatziolos M. 2008.** Coral reefs under rapid climate change and ocean acidification. *Science* **318**:1737–1742 DOI [10.1126/science.1152509](https://doi.org/10.1126/science.1152509).
- Hoogenboom MO, Frank GE, Chase TJ, Jurriaans S, Álvarez Noriega M, Peterson K, Critchell K, Berry KLE, Nicolet KJ, Ramsby B, Paley AS. 2017.** Environmental drivers of variation in bleaching severity of *Acropora* Species during an extreme thermal anomaly [Original Research]. *Frontiers in Marine Science* **4(376)**:1–16 DOI [10.3389/fmars.2017.00376](https://doi.org/10.3389/fmars.2017.00376).
- Jones AM, Berkelmans R. 2011.** Tradeoffs to thermal acclimation: energetics and reproduction of a reef coral with heat tolerant *Symbiodinium* Type-D. *Journal of Marine Biology* **2011**:185890 DOI [10.1155/2011/185890](https://doi.org/10.1155/2011/185890).
- Kemp DW, Hernandez-Pech X, Iglesias-Prieto R, WK Fitt, Schmidt GW. 2014.** Community dynamics and physiology of *Symbiodinium* spp. before, during, and after a coral bleaching event. *Limnology and Oceanography* **59(3)**:788–797 DOI [10.4319/lo.2014.59.3.0788](https://doi.org/10.4319/lo.2014.59.3.0788).
- Keshavmurthy S, Tee HS, Kao KW, Wang JT, Chen CA. 2020.** Specificity trumps flexibility-location-based stable associations between Symbiodiniaceae genera and *Platygyra verweyi* (Scleractinia; Merulinidae). *PeerJ* **8**:e8791 DOI [10.7717/peerj.8791](https://doi.org/10.7717/peerj.8791).
- Klepac C, Beal J, Kenkel C, Sproles A, Polinski J, Williams M, Matz M, Voss J. 2015.** Seasonal stability of coral-*Symbiodinium* associations in the subtropical coral habitat of St. Lucie Reef, Florida. *Marine Ecology Progress Series* **532**:137–151 DOI [10.3354/meps11369](https://doi.org/10.3354/meps11369).
- Kleypas J, McManus J, MeÑEz L. 1999.** Environmental limits to coral reef development: where do we draw the line? *American Zoologist* **39**:146–159 DOI [10.1093/icb/39.1.146](https://doi.org/10.1093/icb/39.1.146).
- Kuanui P, Chavanich S, Raksasab C, Viyakarn V. 2008.** Lunar periodicity of larval release and larval development of *Pocillopora damicornis* in Thailand. *Marine & Freshwater Research* **11**:375–377.
- Kuanui P, Chavanich S, Viyakarn V, Omori M, Fujita T, Lin C. 2020.** Effect of light intensity on survival and photosynthetic efficiency of cultured corals of different ages. *Estuarine, Coastal and Shelf Science* **235**:106515 DOI [10.1016/j.ecss.2019.106515](https://doi.org/10.1016/j.ecss.2019.106515).
- Kuanui P, Chavanich S, Viyakarn V, Omori M, Lin C. 2015.** Effects of temperature and salinity on survival rate of cultured corals and photosynthetic efficiency of zooxanthellae in coral tissues. *Ocean Science Journal* **50**:263–268 DOI [10.1007/s12601-015-0023-3](https://doi.org/10.1007/s12601-015-0023-3).
- Ladrière O, Penin L, Van Lierde E, Vidal-Dupiol J, Kayal M, Roberty S, Poulicek M, Adjeroud M. 2014.** Natural spatial variability of algal endosymbiont density in the coral *Acropora globiceps*: a small-scale approach along environmental gradients around Moorea (French Polynesia). *Journal of the Marine Biological Association of the United Kingdom* **94(1)**:65–74 DOI [10.1017/S0025315413001252](https://doi.org/10.1017/S0025315413001252).

- LaJeunesse TC. 2002.** Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. *Marine Biology* **141**(2):387–400
DOI [10.1007/s00227-002-0829-2](https://doi.org/10.1007/s00227-002-0829-2).
- LaJeunesse T, Bhagooli R, Hidaka M, Devantier L, Done T, Schmidt G, Fitt W, Hoegh-Guldberg O. 2004.** Closely related *Symbiodinium* spp. differ in relative dominance in coral reef host communities across environmental, latitudinal and biogeographic gradients. *Marine Ecology Progress Series* **284**:147–161 DOI [10.3354/meps284147](https://doi.org/10.3354/meps284147).
- LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, Santos SR. 2018.** Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Current Biology* **28**(16):2570–2580
DOI [10.1016/j.cub.2018.07.008](https://doi.org/10.1016/j.cub.2018.07.008).
- LaJeunesse TC, Pettay D, Sampayo E, Phongsuwan N, Brown B, Obura D, Hoegh-Guldberg O, Fitt W. 2010.** Long-standing environmental conditions, geographic isolation and host-symbiont specificity influence the relative ecological dominance and genetic diversification of coral endosymbionts in the genus *Symbiodinium*. *Journal of Biogeography* **37**:785–800 DOI [10.1111/j.1365-2699.2010.02273.x](https://doi.org/10.1111/j.1365-2699.2010.02273.x).
- Levas SJ, Grottoli AG, Hughes A, Osburn CL, Matsui Y. 2013.** Physiological and biogeochemical traits of bleaching and recovery in the mounding species of coral *Porites lobata*: Implications for resilience in mounding corals. *PLOS ONE* **8**(5):e63267
DOI [10.1371/journal.pone.0063267](https://doi.org/10.1371/journal.pone.0063267).
- Lewis CL, Coffroth MA. 2004.** The acquisition of exogenous algal symbionts by an octocoral after bleaching. *Science* **304**(5676):1490–1492 DOI [10.1126/science.1097323](https://doi.org/10.1126/science.1097323).
- Li S, Yu K, Shi Q, Chen T, Zhao M, Zhao J. 2008.** Interspecies and spatial diversity in the symbiotic zooxanthellae density in corals from northern South China Sea and its relationship to coral reef bleaching. *Chinese Science Bulletin* **53**(2):295–303
DOI [10.1007/s11434-007-0514-4](https://doi.org/10.1007/s11434-007-0514-4).
- Little AF, Van Oppen MJH, Willis BL. 2004.** Flexibility in algal endosymbioses shapes growth in reef corals. *Science* **304**(5676):1492–1494 DOI [10.1126/science.1095733](https://doi.org/10.1126/science.1095733).
- Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H, Van Woesik R. 2001.** Coral bleaching: the winners and the losers. *Ecology Letters* **4**(2):122–131
DOI [10.1046/j.1461-0248.2001.00203.x](https://doi.org/10.1046/j.1461-0248.2001.00203.x).
- McCowan DM, Pratchett M, Paely AS, Seeley M, Baird A. 2011.** A comparison of two methods of obtaining densities of zooxanthellae in *Acropora millepora*. *Galaxea Journal of Coral Reef Studies* **13**:1–6 DOI [10.3755/galaxea.13.29](https://doi.org/10.3755/galaxea.13.29).
- McMurdie PJ, Holmes S. 2013.** phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE* **8**(4):e61217
DOI [10.1371/journal.pone.0061217](https://doi.org/10.1371/journal.pone.0061217).
- Muller-Parker G, Davy SK. 2001.** Temperate and tropical algal-sea anemone symbioses. *Invertebrate Biology* **120**(2):104–123 DOI [10.1111/j.1744-7410.2001.tb00115.x](https://doi.org/10.1111/j.1744-7410.2001.tb00115.x).
- Ng TY, Ang P. 2016.** Low symbiont diversity as a potential adaptive strategy in a marginal non-reefal environment: a case study of corals in Hong Kong. *Coral Reefs* **35**(3):941–957 DOI [10.1007/s00338-016-1458-4](https://doi.org/10.1007/s00338-016-1458-4).

- Nielsen DA, Petrou K, Gates RD. 2018. Coral bleaching from a single cell perspective. *The ISME Journal* 12(6):1558–1567 DOI 10.1038/s41396-018-0080-6.
- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin P, O'Hara RB, Simpson G, Solymos P, Stevens MHH, Szöcs E, Wagner H. 2020. Vegan community ecology package version 2.5-7 2020. Available at www.researchgate.net/publication/346579465 (accessed on 12 January 2020).
- Peng SE, Chen CS, Song YF, Huang HT, Jiang PL, Chen WN, Fang LS, Lee YC. 2012. Assessment of metabolic modulation in free-living versus endosymbiotic *Symbiodinium* using synchrotron radiation-based infrared microspectroscopy. *Biology Letters* 8(3):434–437 DOI 10.1098/rsbl.2011.0893.
- Pierangelini M, Thiry M, Cardol P. 2020. Different levels of energetic coupling between photosynthesis and respiration do not determine the occurrence of adaptive responses of Symbiodiniaceae to global warming. *New Phytologist* 228(3):855–868 DOI 10.1111/nph.16738.
- Pillay R, Willis B, Terashima H. 2005. Trends in the density of zooxanthellae in *Acropora millepora* (Ehrenberg, 1834) at the Palm Island Group, Great Barrier Reef, Australia. *Symbiosis* 38:209–226.
- Pootakham W, Mhuantong W, Putchim L, Yoocha T, Sonthirod C, Kongkachana W, Sangsrakru D, Naktang C, Jomchai N, Thongtham N, Tangphatsornruang S. 2018. Dynamics of coral-associated microbiomes during a thermal bleaching event. *Microbiologyopen* 7(5):e00604 DOI 10.1002/mbo3.604.
- Pootakham W, Mhuantong W, Yoocha T, Sangsrakru D, Kongkachana W, Sonthirod C, Naktang C, Jomchai N, Yeemin TSUT, Pongsakun S, Sutthacheep M, Tangphatsornruang S. 2021. Taxonomic profiling of Symbiodiniaceae and bacterial communities associated with Indo-Pacific corals in the Gulf of Thailand using PacBio sequencing of full-length ITS and 16S rRNA genes. *Genomics* 113(4):2717–2729 DOI 10.1016/j.ygeno.2021.06.001.
- Poquita-Du RC, Huang D, Chou LM, Todd PA. 2020. The contribution of stress-tolerant endosymbiotic dinoflagellate *Durusdinium* to *Pocillopora acuta* survival in a highly urbanized reef system. *Coral Reefs* 39(3):745–755 DOI 10.1007/s00338-020-01902-0.
- Putchim L, Phongsuwan N, Yaemarunpattana C, Thongtham N, Richter C. 2017. Long-term changes in the susceptibility of corals to thermal stress around Phuket, Thailand. *PeerJ Preprints* 5:e2979v1 DOI 10.7287/peerj.preprints.2979v1.
- Putnam HM, Stat M, Pochon X, Gates RD. 2012. Endosymbiotic flexibility associates with environmental sensitivity in scleractinian corals. *Proceedings. Biological Sciences* 279(1746):4352–4361 DOI 10.1098/rspb.2012.1454.
- Qin Z, Yu K, Chen B, Wang Y, Liang J, Luo W, Xu L, Huang X. 2019. Diversity of Symbiodiniaceae in 15 coral species from the Southern South China Sea: potential relationship with coral thermal adaptability. *Frontiers in Microbiology* 10:1–12 DOI 10.3389/fmicb.2019.02343.
- Quigley KM, Warner PA, Bay LK, Willis BL. 2018. Unexpected mixed-mode transmission and moderate genetic regulation of *Symbiodinium* communities in a brooding coral. *Heredity* 121(6):524–536 DOI 10.1038/s41437-018-0059-0.

- Quigley KM, Willis BL, Bay LK. 2016.** Maternal effects and *Symbiodinium* community composition drive differential patterns in juvenile survival in the coral *Acropora tenuis*. *Royal Society Open Science* **3**(10):160471 DOI [10.1098/rsos.160471](https://doi.org/10.1098/rsos.160471).
- Quigley KM, Willis BL, Bay LK. 2017.** Heritability of the *Symbiodinium* community in vertically- and horizontally-transmitting broadcast spawning corals. *Scientific Reports* **7**(1):8219 DOI [10.1038/s41598-017-08179-4](https://doi.org/10.1038/s41598-017-08179-4).
- Quigley KM, Willis BL, Kenkel CD. 2019.** Transgenerational inheritance of shuffled symbiont communities in the coral *Montipora digitata*. *Scientific Reports* **9**(1):13328 DOI [10.1038/s41598-019-50045-y](https://doi.org/10.1038/s41598-019-50045-y).
- Reimer J. 2007.** Seasonal changes in the morphological condition of symbiotic dinoflagellates (*Symbiodinium* spp.) of *Zoanthus sansibaricus* (Anthozoa: Hexacorallia) over a latitudinal range in southern Japan. *South Pacific Study* **27**:1–24.
- Richmond R, Hunter C. 1990.** Reproduction and recruitment of corals: comparisons among the Caribbean, the tropical Pacific, and the Red Sea. *Marine Ecology Progress Series* **60**:185–203 DOI [10.3354/meps060185](https://doi.org/10.3354/meps060185).
- Rodriguez-Lanetty M, Loh W, Carter D, Hoegh-Guldberg O. 2001.** Latitudinal variability in symbiont specificity within the widespread scleractinian coral *Plesiastrea versipora*. *Marine Biology* **138**(6):1175–1181 DOI [10.1007/s002270100536](https://doi.org/10.1007/s002270100536).
- Rouze H, Lecellier G, Pochon X, Torda G, Berteaux-Lecellier V. 2019.** Unique quantitative Symbiodiniaceae signature of coral colonies revealed through spatio-temporal survey in Moorea. *Scientific Reports* **9**(1):7921 DOI [10.1038/s41598-019-44017-5](https://doi.org/10.1038/s41598-019-44017-5).
- Rouze H, Lecellier G, Saulnier D, Berteaux-Lecellier V. 2016.** Symbiodinium clades A and D differentially predispose *Acropora cytherea* to disease and *Vibrio* spp. colonization. *Ecology and Evolution* **6**(2):560–572 DOI [10.1002/ece3.1895](https://doi.org/10.1002/ece3.1895).
- Rouze H, Lecellier GJ, Saulnier D, Planes S, Gueguen Y, Wirshing HH, Berteaux-Lecellier V. 2017.** An updated assessment of *Symbiodinium* spp. that associate with common scleractinian corals from Moorea (French Polynesia) reveals high diversity among background symbionts and a novel finding of clade B. *PeerJ* **5**:e2856 DOI [10.7717/peerj.2856](https://doi.org/10.7717/peerj.2856).
- Rowan R. 2004.** Thermal adaptation in reef coral symbionts. *Nature* **430**(7001):742–742 DOI [10.1038/430742a](https://doi.org/10.1038/430742a).
- Sakai K, Singh T, Iguchi A. 2019.** Bleaching and post-bleaching mortality of *Acropora* corals on a heat-susceptible reef in 2016. *PeerJ* **7**:e8138 DOI [10.7717/peerj.8138](https://doi.org/10.7717/peerj.8138).
- Sampayo E, Ridgway T, Bongaerts P, Hoegh-Guldberg O. 2008.** Bleaching susceptibility and mortality of corals are determined by fine-scale differences in symbiont type. *Proceedings of the National Academy of Sciences of the United States of America* **105**:10444–10449 DOI [10.1073/pnas.0708049105](https://doi.org/10.1073/pnas.0708049105).
- Sampayo EM, Ridgway T, Franceschinis L, Roff G, Hoegh-Guldberg O, Dove S. 2016.** Coral symbioses under prolonged environmental change: living near tolerance range limits. *Scientific Reports* **6**(1):36271 DOI [10.1038/srep36271](https://doi.org/10.1038/srep36271).
- Scheufen T, Iglesias-Prieto R, Enríquez S. 2017.** Changes in the number of symbionts and *Symbiodinium* cell pigmentation modulate differentially coral light

- absorption and photosynthetic performance. *Frontiers in Marine Science* 4:309 DOI 10.3389/fmars.2017.00309.
- Schoepf V, Stat M, Falter JL, McCulloch MT. 2015.** Limits to the thermal tolerance of corals adapted to a highly fluctuating, naturally extreme temperature environment. *Scientific Reports* 5(1):17639 DOI 10.1038/srep17639.
- Shinzato C, Zayasu Y, Kanda M, Kawamitsu M, Satoh N, Yamashita H, Suzuki G. 2018.** Using seawater to document coral-zooxanthella diversity: a new approach to coral reef monitoring using environmental DNA [Technology Report]. *Frontiers in Marine Science* 5(28):1–12 DOI 10.3389/fmars.2018.00028.
- Silverstein RN, Correa AM, Baker AC. 2012.** Specificity is rarely absolute in coral-algal symbiosis: implications for coral response to climate change. *Proceedings. Biological Sciences* 279(1738):2609–2618 DOI 10.1098/rspb.2012.0055.
- Silverstein R, Correa A, LaJeunesse T, Baker A. 2011.** Novel algal symbiont (*Symbiodinium* spp.) diversity in reef corals of Western Australia. *Marine Ecology Progress Series* 422:63–75 DOI 10.3354/meps08934.
- Simpson EH. 1949.** Measurement of diversity. *Nature* 163(4148):688–688 DOI 10.1038/163688a0.
- Sproles AE, Oakley CA, Krueger T, Grossman AR, Weis VM, Meibom A, Davy SK. 2020.** Sub-cellular imaging shows reduced photosynthetic carbon and increased nitrogen assimilation by the non-native endosymbiont *Durusinium trenchii* in the model cnidarian *Aiptasia*. *Environmental Microbiology* 22(9):3741–3753 DOI 10.1111/1462-2920.15142.
- Stat M, Loh W, Hoegh-Guldberg O, Carter D. 2008.** Host symbiont acquisition strategy drives *Symbiodinium* diversity in the southern Great Barrier Reef. *Coral Reefs* 27:763–772 DOI 10.1007/s00338-008-0412-5.
- Stat M, Pochon X, Franklin EC, Bruno JF, Casey KS, Selig ER, Gates RD. 2013.** The distribution of the thermally tolerant symbiont lineage (*Symbiodinium* clade D) in corals from Hawaii: correlations with host and the history of ocean thermal stress. *Ecology and Evolution* 3(5):1317–1329 DOI 10.1002/ece3.556.
- Stimson J. 1997.** The annual cycle of density of zooxanthellae in the tissues of field and laboratory-held *Pocillopora damicornis* (Linnaeus). *Journal of Experimental Marine Biology and Ecology* 214(1):35–48 DOI 10.1016/S0022-0981(96)02753-0.
- Suwa R, Hirose M, Hidaka M. 2008.** Seasonal fluctuation in zooxanthella genotype composition and photophysiology in the corals *Pavona divaricata* and *P. decussata*. *Marine Ecology Progress Series* 361:129–137 DOI 10.3354/meps07372.
- Sweet MJ. 2014.** Symbiodinium diversity within *Acropora muricata* and the surrounding environment. *Marine Ecology* 35(3):343–353 DOI 10.1111/maec.12092.
- Tambutté E, Allemand D, Zoccola D, Meibom A, Lotto S, Segonds N, Tambutté S. 2007.** Observations of the tissue-skeleton interface in the scleractinian coral *Stylophora pistillata*. *Coral Reefs* 26:517–529 DOI 10.1007/s00338-007-0263-5.
- Tan Y, Wainwright B, Afiq-Rosli L, Ip Y, Nie L, Nguyen N, Pointing S, Huang D. 2020.** Endosymbiont diversity and community structure in *Porites lutea* from Southeast

- Asia are driven by a suite of environmental variables. *Symbiosis* **80**:269–277
DOI 10.1007/s13199-020-00671-2.
- Terraneo TI, Fusi M, Hume BCC, Arrigoni R, Voolstra CR, Benzoni F, Forsman ZH, Berumen ML. 2019.** Environmental latitudinal gradients and host-specificity shape Symbiodiniaceae distribution in Red Sea *Porites* corals. *Journal of Biogeography* **46**(10):2323–2335 DOI 10.1111/jbi.13672.
- Thinesh T, Meenatchi R, Jose PA, Kiran GS, Selvin J. 2019.** Differential bleaching and recovery pattern of southeast Indian coral reef to 2016 global mass bleaching event: occurrence of stress-tolerant symbiont *Durussdinium* (Clade D) in corals of Palk Bay. *Marine Pollution Bulletin* **145**:287–294 DOI 10.1016/j.marpolbul.2019.05.033.
- Thornhill D, Fitt W, Schmidt G. 2006.** Highly stable symbioses among western Atlantic brooding corals. *Coral Reefs* **25**:515–519 DOI 10.1007/s00338-006-0157-y.
- Thornhill DJ, Howells EJ, Wham DC, Steury TD, Santos SR. 2017.** Population genetics of reef coral endosymbionts (Symbiodinium, Dinophyceae). *Molecular Ecology* **26**(10):2640–2659 DOI 10.1111/mec.14055.
- Thornhill DJ, Lewis AM, Wham DC, LaJeunesse TC. 2014.** Host-speciast lineages dominate the adaptive radiation of reef coral endosymbionts. *Evolution* **68**(2):352–367 DOI 10.1111/evo.12270.
- Thornhill DJ, Xiang Y, Fitt WK, Santos SR. 2009.** Reef endemism, host specificity and temporal stability in populations of symbiotic dinoflagellates from two ecologically dominant Caribbean corals. *PLOS ONE* **4**(7):e6262
DOI 10.1371/journal.pone.0006262.
- Ulstrup KE, Van Oppen MJ. 2003.** Geographic and habitat partitioning of genetically distinct zooxanthellae (*Symbiodinium*) in *Acropora* corals on the Great Barrier Reef. *Molecular Ecology* **12**(12):3477–3484 DOI 10.1046/j.1365-294x.2003.01988.x.
- Van Oppen M, Mieog J, Sanchez C, Fabricius K. 2005.** Diversity of algal endosymbionts (zooxanthellae) in octocorals: the roles of geography and host relationships. *Molecular Ecology* **14**:2403–2417 DOI 10.1111/j.1365-294X.2005.02545.x.
- Vidal-Dupiol J, Adjeroud M, Roger E, Foure L, Duval D, Mone Y, Ferrier-Pages C, Tambutte E, Tambutte S, Zoccola D, Allemand D, Mitta G. 2009.** Coral bleaching under thermal stress: putative involvement of host/symbiont recognition mechanisms. *BMC Physiology* **9**:14 DOI 10.1186/1472-6793-9-14.
- Viyakarn V, Chavanich S, Raksasab C, Loyjiw T. 2009.** New coral community on a breakwater in Thailand. *Coral Reefs* **28**:427–427 DOI 10.1007/s00338-008-0453-9.
- Wall CB, Kaluhiokalani M, Popp BN, Donahue MJ, Gates RD. 2020.** Divergent symbiont communities determine the physiology and nutrition of a reef coral across a light-availability gradient. *The ISME Journal* **14**(4):945–958
DOI 10.1038/s41396-019-0570-1.
- Wang JT, Keshavmurthy S, Chu TY, Chen CA. 2017.** Diverse responses of Symbiodinium types to menthol and DCMU treatment. *PeerJ* **5**:e3843 DOI 10.7717/peerj.3843.
- Weis VM. 2008.** Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of symbiosis. *Journal of Experimental Biology* **211**(Pt 19):3059–3066
DOI 10.1242/jeb.009597.

- Wham DC, Ning G, LaJeunesse TC. 2017.** Symbiodinium glynnii sp. nov. a species of stress-tolerant symbiotic dinoflagellates from pocilloporid and montiporid corals in the Pacific Ocean. *Phycologia* **56**(4):396–409 DOI [10.2216/16-86.1](https://doi.org/10.2216/16-86.1).
- Wilkinson S, Fisher P, Van Oppen M, Davy S. 2015.** Intra-genomic variation in symbiotic dinoflagellates: recent divergence or recombination between lineages? *BMC Evolutionary Biology* **15**:46 DOI [10.1186/s12862-015-0325-1](https://doi.org/10.1186/s12862-015-0325-1).
- Xu L, Yu K-F, Li S, Liu G, Tao S, Shi Q, Chen T, Zhang H. 2017.** Interseasonal and interspecies diversities of *Symbiodinium* density and effective photochemical efficiency in five dominant reef coral species from Luhuitou fringing reef, northern South China Sea. *Coral Reefs* **36**:477–487 DOI [10.1007/s00338-016-1532-y](https://doi.org/10.1007/s00338-016-1532-y).
- Yamashita H, Suzuki G, Kai S, Hayashibara T, Koike K. 2014.** Establishment of coral–algal symbiosis requires attraction and selection. *PLOS ONE* **9**(5):e97003 DOI [10.1371/journal.pone.0097003](https://doi.org/10.1371/journal.pone.0097003).
- Yuyama I, Higuchi T. 2014.** Comparing the effects of symbiotic algae (*Symbiodinium*) clades C1 and D on early growth stages of *Acropora tenuis*. *PLOS ONE* **9**(6):e98999 DOI [10.1371/journal.pone.0098999](https://doi.org/10.1371/journal.pone.0098999).
- Yuyama I, Ishikawa M, Nozawa M, Yoshida M-a, Ikeo K. 2018.** Transcriptomic changes with increasing algal symbiont reveal the detailed process underlying establishment of coral–algal symbiosis. *Scientific Reports* **8**(1):16802 DOI [10.1038/s41598-018-34575-5](https://doi.org/10.1038/s41598-018-34575-5).
- Yuyama I, Nakamura T, Higuchi T, Hidaka M. 2016.** Different stress tolerances of juveniles of the coral *Acropora tenuis* associated with clades C1 and D *Symbiodinium*. *Zoological Studies* **55**:e19 DOI [10.6620/ZS.2016.55-19](https://doi.org/10.6620/ZS.2016.55-19).
- Zhou J, Huang H, Beardall J, Gao K. 2017.** Effect of UV radiation on the expulsion of *Symbiodinium* from the coral *Pocillopora damicornis*. *Journal of Photochemistry and Photobiology B* **166**:12–17 DOI [10.1016/j.jphotobiol.2016.11.003](https://doi.org/10.1016/j.jphotobiol.2016.11.003).
- Ziegler M, Stone E, Colman D, Takacs-Vesbach C, Shepherd U. 2018.** Patterns of *Symbiodinium* (Dinophyceae) diversity and assemblages among diverse hosts and the coral reef environment of Lizard Island, Australia. *Journal of Phycology* **54**(4):447–460 DOI [10.1111/jpy.12749](https://doi.org/10.1111/jpy.12749).